arXiv:1201.4406v1 [math-ph] 20 Jan 2012

Fundamental solution of the Laplacian in the hyperboloid model of hyperbolic geometry

H S Cohl[1,2] **and E G Kalnins**[3]

[1]Information Technology Laboratory, National Institute of Standards and Technology, Gaithersburg, MD, USA
[2]Department of Mathematics, University of Auckland, Auckland, New Zealand
[3]Department of Mathematics, University of Waikato, Hamilton, New Zealand

E-mail: hcohl@nist.gov

Abstract. Due to the isotropy of d-dimensional hyperbolic space, one expects there to exist a spherically symmetric fundamental solution for its corresponding Laplace-Beltrami operator. The R-radius hyperboloid model of hyperbolic geometry $\mathbf{H}_R^d$ with $R > 0$, represents a Riemannian manifold with negative-constant sectional curvature. We obtain a spherically symmetric fundamental solution of Laplace's equation on this manifold in terms of its geodesic radius. We give several matching expressions for this fundamental solution including a definite integral over reciprocal powers of the hyperbolic sine, finite summation expression over hyperbolic functions, Gauss hypergeometric functions, and in terms of the associated Legendre function of the second kind with order and degree given by $d/2 - 1$ with real argument greater than unity. We also demonstrate uniqueness for a fundamental solution of Laplace's equation on this manifold in terms of a vanishing decay at infinity.

PACS numbers: 02.30.Em, 02.30.Gp, 02.30.Jr, 02.40.Ky

AMS classification scheme numbers: 31C12, 32Q45, 33C05, 35A08, 35J05

1. Introduction

We compute closed-form expressions of a spherically symmetric Green's function (fundamental solution) of the Laplacian (Laplace-Beltrami operator) on a Riemannian manifold of constant negative sectional curvature, namely the hyperboloid model of hyperbolic geometry. Useful background material relevant for this paper can be found in Vilenkin (1968) [25], Thurston (1997) [23], Lee (1997) [18] and Pogosyan & Winternitz (2002) [22].

This paper is organized as follows. In section 2 we describe the hyperboloid model of hyperbolic geometry and its corresponding metric, global geodesic distance function, Laplacian and geodesic polar coordinate systems which parametrize points in this model. In section 3 for the hyperboloid model of hyperbolic geometry, we show how to compute radial harmonics in a geodesic polar coordinate system and derive several alternative expressions for a radial fundamental solution of the Laplacian on the d-dimensional R-radius hyperboloid with $R > 0$. In section 4 we prove that our derived fundamental solution is unique in terms of a vanishing decay at infinity.

Throughout this paper we rely on the following definitions. For $a_1, a_2, \ldots \in \mathbf{C}$, if $i, j \in \mathbf{Z}$ and $j < i$ then $\sum_{n=i}^{j} a_n = 0$ and $\prod_{n=i}^{j} a_n = 1$. The set of natural numbers is given by $\mathbf{N} := \{1, 2, \ldots\}$, the set $\mathbf{N}_0 := \{0, 1, 2, \ldots\} = \mathbf{N} \cup \{0\}$, and the set $\mathbf{Z} := \{0, \pm 1, \pm 2, \ldots\}$. The set $\mathbf{R}$ represents the real numbers.

2. The hyperboloid model of hyperbolic geometry

Hyperbolic space in d-dimensions is a fundamental example of a space exhibiting hyperbolic geometry. It was developed independently by Lobachevsky and Bolyai around 1830 (see Trudeau (1987) [24]). It is a geometry analogous to Euclidean geometry, but such that Euclid's parallel postulate is no longer assumed to hold.

There are several models of d-dimensional hyperbolic geometry including the Klein (see Figure 1), Poincaré (see Figure 2), hyperboloid, upper-half space and hemisphere models (see Thurston (1997) [23]). The hyperboloid model for d-dimensional hyperbolic space is closely related to the Klein and Poincaré models: each can be obtained projectively from the others. The upper-half space and hemisphere models can be obtained from one another by inversions with the Poincaré model (see section 2.2 in Thurston (1997) [23]). The model we will be focusing on in this paper is the hyperboloid model.

The hyperboloid model, also known as the Minkowski or Lorentz models, are models of d-dimensional hyperbolic geometry in which points are represented by the upper sheet (submanifold) S^+ of a two-sheeted hyperboloid embedded in the Minkowski space $\mathbf{R}^{d,1}$. Minkowski space is a $(d+1)$-dimensional pseudo-Riemannian manifold which is a real finite-dimensional vector space, with coordinates given by $\mathbf{x} = (x_0, x_1, \ldots, x_d)$. It is equipped with a nondegenerate, symmetric bilinear form, the Minkowski bilinear form

$$[\mathbf{x}, \mathbf{y}] = x_0 y_0 - x_1 y_1 - \ldots - x_d y_d.$$



Figure 1. This figure is a graphical depiction of stereographic projection from the hyperboloid model to the Klein model of hyperbolic space.

The above bilinear form is symmetric, but not positive-definite, so it is not an inner product. It is defined analogously with the Euclidean inner product for $\mathbf{R}^{d+1}$

$$(\mathbf{x}, \mathbf{y}) = x_0 y_0 + x_1 y_1 + \ldots + x_d y_d.$$

The variety $[\mathbf{x}, \mathbf{x}] = x_0^2 - x_1^2 - \ldots - x_d^2 = R^2$, for $\mathbf{x} \in \mathbf{R}^{d,1}$, using the language of Beltrami (1869) [3] (see also p. 504 in Vilenkin (1968) [25]), defines a pseudo-sphere of radius R. Points on the pseudo-sphere with zero radius coincide with a cone. Points on the pseudo-sphere with radius greater than zero lie within this cone, and points on the pseudo-sphere with purely imaginary radius lie outside the cone.

For $R \in (0, \infty)$, we refer to the variety $[\mathbf{x}, \mathbf{x}] = R^2$ as the R-radius hyperboloid $\mathbf{H}_R^d$.



Figure 2. This figure is a graphical depiction of stereographic projection from the hyperboloid model to the Poincaré model of hyperbolic space.

This variety is a maximally symmetric, simply connected, d-dimensional Riemannian manifold with negative-constant sectional curvature (given by $-1/R^2$, see for instance p. 148 in Lee (1997) [18]), whereas Euclidean space $\mathbf{R}^d$ equipped with the Pythagorean norm, is a space with zero sectional curvature. For a fixed $R \in (0, \infty)$, the R-radius hypersphere $\mathbf{S}^d_R$, is an example of a space (submanifold) with positive constant sectional curvature (given by $1/R^2$). We denote unit radius hyperboloid by $\mathbf{H}^d := \mathbf{H}^d_1$ and the unit radius hypersphere by $\mathbf{S}^d := \mathbf{S}^d_1$.

In our discussion of a fundamental solution for the Laplacian in the hyperboloid model of hyperbolic geometry, we focus on the positive radius pseudo-sphere which can

be parametrized through *subgroup-type coordinates,* i.e. those which correspond to a maximal subgroup chain $O(d,1) \supset \ldots$ (see for instance Pogosyan & Winternitz (2002) [22]). There exist separable coordinate systems which parametrize points on the positive radius pseudo-sphere (i.e. such as those which are analogous to parabolic coordinates, etc.) which can not be constructed using maximal subgroup chains (we will no longer discuss these).

Geodesic polar coordinates are coordinates which correspond to the maximal subgroup chain given by $O(d,1) \supset O(d) \supset \ldots$. What we will refer to as *standard geodesic polar coordinates* correspond to the subgroup chain given by $O(d,1) \supset O(d) \supset O(d-1) \supset \cdots \supset O(2)$. Standard geodesic polar coordinates (see Olevskiĭ (1950) [20]; Grosche, Pogosyan & Sissakian (1997) [13]), similar to standard hyperspherical coordinates in Euclidean space, can be given by

$$\left.\begin{aligned} x_0 &= R\cosh r \\ x_1 &= R\sinh r\cos\theta_1 \\ x_2 &= R\sinh r\sin\theta_1\cos\theta_2 \\ &\vdots \\ x_{d-2} &= R\sinh r\sin\theta_1\cdots\cos\theta_{d-2} \\ x_{d-1} &= R\sinh r\sin\theta_1\cdots\sin\theta_{d-2}\cos\phi \\ x_d &= R\sinh r\sin\theta_1\cdots\sin\theta_{d-2}\sin\phi, \end{aligned}\right\} \tag{1}$$

where $r \in [0,\infty)$, $\phi \in [0,2\pi)$, and $\theta_i \in [0,\pi]$ for $i \in \{1,\ldots,d-2\}$.

The isometry group of the space $\mathbf{H}^d_R$ is the pseudo-orthogonal group $SO(d,1)$, the Lorentz group in $(d+1)$-dimensions. Hyperbolic space $\mathbf{H}^d_R$, can be identified with the quotient space $SO(d,1)/SO(d)$. The isometry group acts transitively on $\mathbf{H}^d_R$. That is, any point on the hyperboloid can be carried, with the help of a Euclidean rotation of $SO(d-1)$, to the point $(\cosh\alpha, \sinh\alpha, 0, \ldots, 0)$, and a hyperbolic rotation

$$\left.\begin{aligned} x'_0 &= -x_1\sinh\alpha + x_0\cosh\alpha \\ x'_1 &= -x_1\cosh\alpha - x_0\sinh\alpha \end{aligned}\right\}$$

maps that point to the origin $(1,0,\ldots,0)$ of the space. In order to study a fundamental solution of Laplace's equation on the hyperboloid, we need to describe how one computes distances in this space.

One may naturally compare distances on the positive radius pseudo-sphere through analogy with the R-radius hypersphere. Distances on the hypersphere are simply given by arc lengths, angles between two arbitrary vectors, from the origin, in the ambient Euclidean space. We consider the d-dimensional hypersphere embedded in $\mathbf{R}^{d+1}$. Points on the hypersphere can be parametrized using hyperspherical coordinate systems. Any parametrization of the hypersphere $\mathbf{S}^d_R$, must have $(\mathbf{x},\mathbf{x}) = x_0^2 + \ldots + x_d^2 = R^2$, with $R > 0$. The distance between two points on the hypersphere $\mathbf{x}, \mathbf{x}' \in \mathbf{S}^d_R$ is given by

$$d(\mathbf{x},\mathbf{x}') = R\gamma = R\cos^{-1}\left(\frac{(\mathbf{x},\mathbf{x}')}{(\mathbf{x},\mathbf{x})(\mathbf{x}',\mathbf{x}')}\right) = R\cos^{-1}\left(\frac{1}{R^2}(\mathbf{x},\mathbf{x}')\right). \tag{2}$$

This is evident from the fact that the geodesics on $\mathbf{S}^d_R$ are great circles (i.e. intersections of $\mathbf{S}^d_R$ with planes through the origin) with constant speed parametrizations (see p. 82 in Lee (1997) [18]).

Accordingly, we now look at the geodesic distance function on the d-dimensional positive radius pseudo-sphere $\mathbf{H}^d_R$. Distances between two points on the positive radius pseudo-sphere are given by the hyperangle between two arbitrary vectors, from the origin, in the ambient Minkowski space. Any parametrization of the hyperboloid $\mathbf{H}^d_R$, must have $[\mathbf{x}, \mathbf{x}] = R^2$. The geodesic distance between two points $\mathbf{x}, \mathbf{x}' \in \mathbf{H}^d_R$ is given by

$$d(\mathbf{x}, \mathbf{x}') = R \cosh^{-1}\left(\frac{[\mathbf{x}, \mathbf{x}']}{[\mathbf{x}, \mathbf{x}][\mathbf{x}', \mathbf{x}']}\right) = R \cosh^{-1}\left(\frac{1}{R^2}[\mathbf{x}, \mathbf{x}']\right), \tag{3}$$

where the inverse hyperbolic cosine with argument $x \in (1, \infty)$ is given by (see (4.37.19) in Olver *et al.* (2010) [21])

$$\cosh^{-1} x = \log\left(x + \sqrt{x^2 - 1}\right).$$

Geodesics on $\mathbf{H}^d_R$ are great hyperbolas (i.e. intersections of $\mathbf{H}^d_R$ with planes through the origin) with constant speed parametrizations (see p. 84 in Lee (1997) [18]). We also define a global function $\rho : \mathbf{H}^d \times \mathbf{H}^d \to [0, \infty)$ which represents the projection of global geodesic distance function (3) on $\mathbf{H}^d_R$ onto the corresponding unit radius hyperboloid $\mathbf{H}^d$, namely

$$\rho(\widehat{\mathbf{x}}, \widehat{\mathbf{x}}') := d(\mathbf{x}, \mathbf{x}')/R, \tag{4}$$

where $\widehat{\mathbf{x}} = \mathbf{x}/R$ and $\widehat{\mathbf{x}}' = \mathbf{x}'/R$.

2.1. The Laplacian on the hyperboloid model

Parametrizations of a submanifold embedded in either a Euclidean or Minkowski space is given in terms of coordinate systems whose coordinates are curvilinear. These are coordinates based on some transformation that converts the standard Cartesian coordinates in the ambient space to a coordinate system with the same number of coordinates as the dimension of the submanifold in which the coordinate lines are curved.

On a d-dimensional Riemannian manifold M (a manifold together with a Riemannian metric g), the Laplace-Beltrami operator (Laplacian) $\Delta : C^p(M) \to C^{p-2}(M)$, $p \geq 2$, in curvilinear coordinates $\xi = (\xi^1, \ldots, \xi^d)$ is given by

$$\Delta = \sum_{i,j=1}^{d} \frac{1}{\sqrt{|g|}} \frac{\partial}{\partial \xi^i}\left(\sqrt{|g|} g^{ij} \frac{\partial}{\partial \xi^j}\right), \tag{5}$$

where $|g| = |\det(g_{ij})|$, the infinitesimal distance is given by

$$ds^2 = \sum_{i,j=1}^{d} g_{ij} d\xi^i d\xi^j, \tag{6}$$

and

$$\sum_{i=1}^{d} g_{ki} g^{ij} = \delta_k^j,$$

where $\delta_i^j \in \{0, 1\}$ is the Kronecker delta defined for all $i, j \in \mathbf{Z}$ such that

$$\delta_i^j := \begin{cases} 1 & \text{if } i = j, \\ 0 & \text{if } i \neq j. \end{cases} \tag{7}$$

For a Riemannian submanifold, the relation between the metric tensor in the ambient space and g_{ij} of (5) and (6) is

$$g_{ij}(\xi) = \sum_{k,l=0}^{d} G_{kl} \frac{\partial x^k}{\partial \xi^i} \frac{\partial x^l}{\partial \xi^j}.$$

On $\mathbf{H}_R^d$ the ambient space is Minkowski, and therefore $G_{ij} = \mathrm{diag}(1, -1, \ldots, -1)$.

The set of all geodesic polar coordinate systems on the hyperboloid correspond to the many ways one can put coordinates on a hyperbolic hypersphere, i.e., the Riemannian submanifold $U \subset \mathbf{H}_R^d$ defined for a fixed $\mathbf{x}' \in \mathbf{H}_R^d$ such that $d(\mathbf{x}, \mathbf{x}') = b = const$, where $b \in (0, \infty)$. These are coordinate systems which correspond to subgroup chains starting with $O(d, 1) \supset O(d) \supset \cdots$, with standard geodesic polar coordinates given by (1) being only one of them. (For a thorough description of these see section X.5 in Vilenkin (1968) [25].) They all share the property that they are described by $(d + 1)$-variables: $r \in [0, \infty)$ plus d-angles each being given by the values $[0, 2\pi)$, $[0, \pi]$, $[-\pi/2, \pi/2]$ or $[0, \pi/2]$ (see Izmest'ev *et al.* (1999, 2001) [16, 17]).

In any of the geodesic polar coordinate systems, the global geodesic distance between any two points on the hyperboloid is given by (cf. (3))

$$d(\mathbf{x}, \mathbf{x}') = R \cosh^{-1}(\cosh r \cosh r' - \sinh r \sinh r' \cos \gamma), \tag{8}$$

where γ is the unique separation angle given in each hyperspherical coordinate system. For instance, the separation angle in standard geodesic polar coordinates (1) is given by the formula

$$\cos\, \gamma = \cos(\phi - \phi') \prod_{i=1}^{d-2} \sin\theta_i \sin\theta_i{}' + \sum_{i=1}^{d-2} \cos\theta_i \cos\theta_i{}' \prod_{j=1}^{i-1} \sin\theta_j \sin\theta_j{}'. \tag{9}$$

Corresponding separation angle formulae for any geodesic polar coordinate system can be computed using (2), (3), and the associated formulae for the appropriate inner-products. Note that by making use of the isometry group $SO(d, 1)$ to map $\mathbf{x}'$ to the origin, then $\rho = Rr$ for $\mathbf{H}_R^d$ and in particular $\rho = r$ for $\mathbf{H}^d$. Hence, for the unit radius hyperboloid, there is no distinction between the global geodesic distance and the r-parameter in a geodesic polar coordinate system. For the R-radius hyperboloid, the only distinction between the global geodesic distance and the r-parameter is the multiplicative constant R.

The infinitesimal distance in a geodesic polar coordinate system on this submanifold is given by

$$ds^2 = R^2(dr^2 + \sinh^2 r \; d\gamma^2), \tag{10}$$

where an appropriate expression for γ in a curvilinear coordinate system is given. If one combines (1), (5), (9) and (10), then in a particular geodesic polar coordinate system, Laplace's equation on $\mathbf{H}^d_R$ is given by

$$\Delta f = \frac{1}{R^2}\left[\frac{\partial^2 f}{\partial r^2} + (d-1)\coth r \frac{\partial f}{\partial r} + \frac{1}{\sinh^2 r}\Delta_{\mathbf{S}^{d-1}} f\right] = 0, \tag{11}$$

where $\Delta_{\mathbf{S}^{d-1}}$ is the corresponding Laplace-Beltrami operator on the unit radius hypersphere $\mathbf{S}^{d-1}$.

3. A Green's function in the hyperboloid model

3.1. Harmonics in geodesic polar coordinates

Geodesic polar coordinate systems partition the R-radius hyperboloid $\mathbf{H}^d_R$ into a family of $(d-1)$-dimensional hyperbolic hyperspheres, each with a radius $r \in (0, \infty)$, on which all possible hyperspherical coordinate systems for $\mathbf{S}^{d-1}$ may be used (see for instance Vilenkin (1968) [25]). One then must also consider the limiting case for $r = 0$ to fill out all of $\mathbf{H}^d_R$. In geodesic polar coordinates one can compute the normalized hyperspherical harmonics in this space by solving the Laplace equation using separation of variables which results in a general procedure which is given explicitly in Izmest'ev *et al.* (1999, 2001) [16, 17]. These angular harmonics are given as general expressions involving trigonometric functions, Gegenbauer polynomials and Jacobi polynomials.

The harmonics in geodesic polar coordinate systems are given in terms of a radial solution multiplied by the angular harmonics. The angular harmonics are eigenfunctions of the Laplace-Beltrami operator on $\mathbf{S}^{d-1}$ with unit radius which satisfy the following eigenvalue problem

$$\Delta_{\mathbf{S}^{d-1}} Y_l^K(\widehat{\mathbf{x}}) = -l(l+d-2)Y_l^K(\widehat{\mathbf{x}}),$$

where $\widehat{\mathbf{x}} \in \mathbf{S}^{d-1}$, $Y_l^K(\widehat{\mathbf{x}})$ are normalized hyperspherical harmonics, $l \in \mathbf{N}_0$ is the angular momentum quantum number, and K stands for the set of $(d-2)$-quantum numbers identifying degenerate harmonics for each l. The degeneracy

$$(2l+d-2)\frac{(d-3+l)!}{l!(d-2)!}$$

(see (9.2.11) in Vilenkin (1968) [25]), tells you how many linearly independent solutions exist for a particular l value and dimension d. The hyperspherical harmonics are normalized such that

$$\int_{\mathbf{S}^{d-1}} Y_l^K(\widehat{\mathbf{x}})\overline{Y_{l'}^{K'}(\widehat{\mathbf{x}})}d\omega = \delta_l^{l'}\delta_K^{K'},$$

where $d\omega$ is the Riemannian (volume) measure (see for instance section 3.4 in Grigor'yan (2009)[12]) on $\mathbf{S}^{d-1}$ which is invariant under the isometry group $SO(d)$ (cf. (12)), and for

$x+iy=z\in\mathbf{C}$, $\overline{z}=x-iy$, represents complex conjugation. The generalized Kronecker delta $\delta_K^{K'}$ (cf. (7)) is defined such that it equals 1 if all of the $(d-2)$-quantum numbers identifying degenerate harmonics for each l coincide, and equals zero otherwise.

Since the angular solutions (hyperspherical harmonics) are well-known (see Chapter IX in Vilenkin (1968) [25]; Chapter 11 in Erdélyi *et al.* (1981) [7]), we will now focus on the radial solutions on $\mathbf{H}_R^d$ in geodesic polar coordinates, which satisfy the following ordinary differential equation (cf. (11)) for all $R\in(0,\infty)$, namely

$$\frac{d^2u}{dr^2}+(d-1)\coth r\frac{du}{dr}-\frac{l(l+d-2)}{\sinh^2 r}u=0.$$

Four solutions to this ordinary differential equation $u_{1\pm}^{d,l},u_{2\pm}^{d,l}:(1,\infty)\to\mathbf{C}$ are given by

$$u_{1\pm}^{d,l}(\cosh r)=\frac{1}{\sinh^{d/2-1}r}P_{d/2-1}^{\pm(d/2-1+l)}(\cosh r),$$

and

$$u_{2\pm}^{d,l}(\cosh r)=\frac{1}{\sinh^{d/2-1}r}Q_{d/2-1}^{\pm(d/2-1+l)}(\cosh r),$$

where $P_\nu^\mu,Q_\nu^\mu:(1,\infty)\to\mathbf{C}$ are associated Legendre functions of the first and second kind respectively (see for instance Chapter 14 in Olver *et al.* (2010) [21]).

Due to the fact that the space $\mathbf{H}_R^d$ is homogeneous with respect to its isometry group, the pseudo-orthogonal group $SO(d,1)$, and therefore an isotropic manifold, we expect that there exist a fundamental solution of Laplace's equation on this space with spherically symmetric dependence. We specifically expect these solutions to be given in terms of associated Legendre functions of the second kind with argument given by $\cosh r$. This associated Legendre function naturally fits our requirements because it is singular at $)r=0$ and vanishes at infinity, whereas the associated Legendre functions of the first kind, with the same argument, are regular at $r=0$ and singular at infinity.

3.2. Fundamental solution of the Laplacian

In computing a fundamental solution of the Laplacian on $\mathbf{H}_R^d$, we know that

$$-\Delta\mathcal{H}_R^d(\mathbf{x},\mathbf{x}')=\delta_g(\mathbf{x},\mathbf{x}'),$$

where g is the Riemannian metric on $\mathbf{H}_R^d$ and $\delta_g(\mathbf{x},\mathbf{x}')$ is the Dirac delta function on the manifold $\mathbf{H}_R^d$. The Dirac delta function is defined for an open set $U\subset\mathbf{H}_R^d$ with $\mathbf{x},\mathbf{x}'\in\mathbf{H}_R^d$ such that

$$\int_U\delta_g(\mathbf{x},\mathbf{x}')d\mathrm{vol}_g=\begin{cases}1 & \text{if } \mathbf{x}'\in U,\\ 0 & \text{if } \mathbf{x}'\notin U,\end{cases}$$

where $d\mathrm{vol}_g$ is the Riemannian (volume) measure, invariant under the isometry group $SO(d,1)$ of the Riemannian manifold $\mathbf{H}_R^d$, given (in standard geodesic polar coordinates) by

$$d\mathrm{vol}_g=R^d\sinh^{d-1}r\,dr\,d\omega:=R^d\sinh^{d-1}r\,dr\,\sin^{d-2}\theta_{d-1}\cdots\sin\theta_2 d\theta_1\cdots d\theta_{d-1}. \tag{12}$$

Notice that as $r \to 0^+$ that $d\mathrm{vol}_g$ goes to the Euclidean measure, invariant under the Euclidean motion group $E(d)$, in spherical coordinates. Therefore in spherical coordinates, we have the following

$$\delta_g(\mathbf{x},\mathbf{x}') = \frac{\delta(r-r')}{R^d \sinh^{d-1} r'} \frac{\delta(\theta_1-\theta_1')\cdots\delta(\theta_{d-1}-\theta_{d-1}')}{\sin\theta_2'\cdots\sin^{d-2}\theta_{d-1}'}.$$

In general since we can add any harmonic function to a fundamental solution of the Laplacian and still have a fundamental solution, we will use this freedom to make our fundamental solution as simple as possible. It is reasonable to expect that there exists a particular spherically symmetric fundamental solution $\mathcal{H}_R^d(\mathbf{x},\mathbf{x}')$ on the hyperboloid with pure radial $\rho(\widehat{\mathbf{x}},\widehat{\mathbf{x}}') := d(\mathbf{x},\mathbf{x}')/R$ (cf. (4)) and constant angular dependence (invariant under rotations centered about the origin), due to the influence of the point-like nature of the Dirac delta function. For a spherically symmetric solution to the Laplace equation, the corresponding $\Delta_{\mathbf{S}^{d-1}}$ term vanishes since only the $l = 0$ term survives. In other words, we expect there to exist a fundamental solution of Laplace's equation such that $\mathcal{H}_R^d(\mathbf{x},\mathbf{x}') = f(\rho)$.

We have proven that on the R-radius hyperboloid $\mathbf{H}_R^d$, a Green's function for the Laplace operator (fundamental solution for the Laplacian) can be given as follows.

Theorem 3.1. *Let $d \in \{2,3,\ldots\}$. Define $\mathcal{I}_d : (0,\infty) \to \mathbf{R}$ as*

$$\mathcal{I}_d(\rho) := \int_\rho^\infty \frac{dx}{\sinh^{d-1} x},$$

$\mathbf{x},\mathbf{x}' \in \mathbf{H}_R^d$, and $\mathcal{H}_R^d : (\mathbf{H}_R^d \times \mathbf{H}_R^d) \setminus \{(\mathbf{x},\mathbf{x}) : \mathbf{x} \in \mathbf{H}_R^d\} \to \mathbf{R}$ defined such that

$$\mathcal{H}_R^d(\mathbf{x},\mathbf{x}') := \frac{\Gamma(d/2)}{2\pi^{d/2}R^{d-2}}\mathcal{I}_d(\rho),$$

where $\rho := \cosh^{-1}([\widehat{\mathbf{x}},\widehat{\mathbf{x}}'])$ is the geodesic distance between $\widehat{\mathbf{x}}$ and $\widehat{\mathbf{x}}'$ on the pseudo-sphere of unit radius $\mathbf{H}^d$, with $\widehat{\mathbf{x}} = \mathbf{x}/R$, $\widehat{\mathbf{x}}' = \mathbf{x}'/R$, then $\mathcal{H}_R^d$ is a fundamental solution for $-\Delta$ where Δ is the Laplace-Beltrami operator on $\mathbf{H}_R^d$. Moreover,

$$\mathcal{I}_d(\rho) = \begin{cases} (-1)^{d/2-1}\dfrac{(d-3)!!}{(d-2)!!}\left[\log\coth\dfrac{\rho}{2} + \cosh\rho\displaystyle\sum_{k=1}^{d/2-1}\frac{(2k-2)!!(-1)^k}{(2k-1)!!\sinh^{2k}\rho}\right] & \text{if } d \text{ even,} \\ \left\{\begin{array}{l} (-1)^{(d-1)/2}\left[\dfrac{(d-3)!!}{(d-2)!!}\right. \\ \qquad\left. + \left(\dfrac{d-3}{2}\right)!\displaystyle\sum_{k=1}^{(d-1)/2}\frac{(-1)^k\coth^{2k-1}\rho}{(2k-1)(k-1)!((d-2k-1)/2)!}\right], \\ \text{or} \\ (-1)^{(d-1)/2}\dfrac{(d-3)!!}{(d-2)!!}\left[1+\cosh\rho\displaystyle\sum_{k=1}^{(d-1)/2}\frac{(2k-3)!!(-1)^k}{(2k-2)!!\sinh^{2k-1}\rho}\right], \end{array}\right\} & \text{if } d \text{ odd.} \end{cases}$$

$$= \frac{1}{(d-1)\cosh^{d-1}\rho}\,{}_2F_1\left(\frac{d-1}{2},\frac{d}{2};\frac{d+1}{2};\frac{1}{\cosh^2\rho}\right),$$

$$= \frac{1}{(d-1)\cosh\rho\,\sinh^{d-2}\rho}\,{}_2F_1\left(\frac{1}{2},1;\frac{d+1}{2};\frac{1}{\cosh^2\rho}\right),$$

$$= \frac{e^{-i\pi(d/2-1)}}{2^{d/2-1}\Gamma\left(d/2\right)\sinh^{d/2-1}\rho}\,Q^{d/2-1}_{d/2-1}(\cosh\rho),$$

where !! *is the double factorial,* ${}_2F_1$ *is the Gauss hypergeometric function, and* Q^{μ}_{ν} *is the associated Legendre function of the second kind.*

In the rest of this section, we develop the material in order to prove this theorem.

Due to the fact that the space $\mathbf{H}^d_R$ is homogeneous with respect to its isometry group $SO(d,1)$, and therefore an isotropic manifold, without loss of generality, we are free to map the point $\mathbf{x}' \in \mathbf{H}^d_R$ to the origin. In this case the global distance function $\rho : \mathbf{H}^d \times \mathbf{H}^d \to [0,\infty)$ coincides with the radial parameter in geodesic polar coordinates, and we may interchange r with ρ accordingly (cf. (8) with $r' = 0$) in our representation of a fundamental solution of Laplace's equation on this manifold. Since a spherically symmetric choice for a fundamental solution of Laplace's equation is harmonic everywhere except at the origin, we may first set $g = f'$ in (11) and solve the first-order equation

$$g' + (d-1)\coth\rho\; g = 0,$$

which is integrable and clearly has the general solution

$$g(\rho) = \frac{df}{d\rho} = c_0\sinh^{1-d}\rho, \tag{13}$$

where $c_0 \in \mathbf{R}$ is a constant which depends on d. Now we integrate (13) to obtain a fundamental solution for the Laplacian in $\mathbf{H}^d_R$

$$\mathcal{H}^d_R(\mathbf{x},\mathbf{x}') = c_0\mathcal{I}_d(\rho) + c_1, \tag{14}$$

where

$$\mathcal{I}_d(\rho) := \int_\rho^\infty \frac{dx}{\sinh^{d-1}x}, \tag{15}$$

and $c_0, c_1 \in \mathbf{R}$ are constants which depend on d. This definite integral result is mentioned in section II.5 of Helgason (1984) [14] and as well in Losev (1986) [19] . Notice that we can add any harmonic function to (14) and still have a fundamental solution of the Laplacian since a fundamental solution of the Laplacian must satisfy

$$\int_{\mathbf{H}^d_R} (-\Delta\varphi)(\mathbf{x}')\mathcal{H}^d_R(\mathbf{x},\mathbf{x}')\; d\mathrm{vol}'_g = \varphi(\mathbf{x}),$$

for all $\varphi \in \mathcal{D}(\mathbf{H}^d_R)$, where $\mathcal{D}$ is the space of test functions, and $d\mathrm{vol}'_g$ is the Riemannian (volume) measure on $\mathbf{H}^d_R$, in the primed coordinates. In particular, we notice that from our definition of $\mathcal{I}_d$ (15) that

$$\lim_{\rho\to\infty}\mathcal{I}_d(\rho) = 0.$$

Therefore it is convenient to set $c_1 = 0$ leaving us with

$$\mathcal{H}_R^d(\mathbf{x}, \mathbf{x}') = c_0 \mathcal{I}_d(\rho). \tag{16}$$

In Euclidean space $\mathbf{R}^d$, a Green's function for Laplace's equation (fundamental solution for the Laplacian) is well-known and is given in the following theorem (see Folland (1976) [8]; p. 94, Gilbarg & Trudinger (1983) [9]; p. 17, Bers *et al.* (1964) [4], p. 211).

Theorem 3.2. *Let $d \in \mathbf{N}$. Define*

$$\mathcal{G}^d(\mathbf{x}, \mathbf{x}') = \begin{cases} \dfrac{\Gamma(d/2)}{2\pi^{d/2}(d-2)} \|\mathbf{x} - \mathbf{x}'\|^{2-d} & \text{if } d = 1 \text{ or } d \geq 3, \\ \dfrac{1}{2\pi} \log \|\mathbf{x} - \mathbf{x}'\|^{-1} & \text{if } d = 2, \end{cases}$$

then $\mathcal{G}_R^d$ is a fundamental solution for $-\Delta$ in Euclidean space $\mathbf{R}^d$, where Δ is the Laplace operator in $\mathbf{R}^d$.

Note most authors only present the above theorem for the case $d \geq 2$ but it is easily-verified to also be valid for the case $d = 1$ as well.

The hyperboloid $\mathbf{H}_R^d$, being a manifold, must behave locally like Euclidean space $\mathbf{R}^d$. Therefore for small ρ we have $e^\rho \simeq 1 + \rho$ and $e^{-\rho} \simeq 1 - \rho$ and in that limiting regime

$$\mathcal{I}_d(\rho) \approx \int_\rho^1 \frac{dx}{x^{d-1}} \simeq \begin{cases} -\log \rho & \text{if } d = 2, \\ \dfrac{1}{\rho^{d-2}} & \text{if } d \geq 3, \end{cases}$$

which has exactly the same singularity as a Euclidean fundamental solution for Laplace's equation. Therefore the proportionality constant c_0 is obtained by matching locally to a Euclidean fundamental solution of Laplace's equation

$$\mathcal{H}_R^d = c_0 \mathcal{I}_d \simeq \mathcal{G}^d,$$

near the singularity located at $\mathbf{x} = \mathbf{x}'$.

We have shown how to compute a fundamental solution of the Laplace-Beltrami operator on the hyperboloid in terms of an improper integral (15). We would now like to express this integral in terms of well-known special functions. A fundamental solution $\mathcal{I}_d$ can be computed using elementary methods through its definition (15). In $d = 2$ we have

$$\mathcal{I}_2(\rho) = \int_\rho^\infty \frac{dx}{\sinh x} = \frac{1}{2} \log \frac{\cosh \rho + 1}{\cosh \rho - 1} = \log \coth \frac{\rho}{2},$$

and in $d = 3$ we have

$$\mathcal{I}_3(\rho) = \int_\rho^\infty \frac{dx}{\sinh^2 x} = \frac{e^{-\rho}}{\sinh \rho} = \coth \rho - 1.$$

This exactly matches up to that given by (3.27) in Hostler (1955) [15]. In $d \in \{4,5,6,7\}$ we have

$$\begin{aligned}
\mathcal{I}_4(\rho) &= -\frac{1}{2}\log\coth\frac{\rho}{2} + \frac{\cosh\rho}{2\sinh^2\rho},\\
\mathcal{I}_5(\rho) &= \frac{1}{3}(\coth^3\rho - 1) - (\coth\rho - 1),\\
\mathcal{I}_6(\rho) &= \frac{3}{8}\log\coth\frac{\rho}{2} + \frac{\cosh\rho}{4\sinh^4\rho} - \frac{3\cosh\rho}{8\sinh^2\rho}, \quad \text{and}\\
\mathcal{I}_7(\rho) &= \frac{1}{5}(\coth^5\rho - 1) - \frac{2}{3}(\coth^3\rho - 1) + \coth\rho - 1.
\end{aligned}$$

Now we prove several equivalent finite summation expressions for $\mathcal{I}_d(\rho)$. We wish to compute the antiderivative $\mathfrak{I}_m : (0,\infty) \to \mathbf{R}$, which is defined as

$$\mathfrak{I}_m(x) := \int \frac{dx}{\sinh^m x},$$

where $m \in \mathbf{N}$. This antiderivative satisfies the following recurrence relation

$$\mathfrak{I}_m(x) = -\frac{\cosh x}{(m-1)\sinh^{m-1} x} - \frac{(m-2)}{(m-1)}\mathfrak{I}_{m-2}(x), \tag{17}$$

which follows from the identity

$$\frac{1}{\sinh^m x} = \frac{\cosh x}{\sinh^m x}\cosh x - \frac{1}{\sinh^{m-2} x},$$

and integration by parts. The antiderivative $\mathfrak{I}_m(x)$ naturally breaks into two separate classes, namely

$$\begin{aligned}
\int \frac{dx}{\sinh^{2n+1} x} &= (-1)^{n+1}\frac{(2n-1)!!}{(2n)!!}\\
&\quad \times \left[\log\coth\frac{x}{2} + \cosh x \sum_{k=1}^{n} \frac{(2k-2)!!(-1)^k}{(2k-1)!!\sinh^{2k} x}\right] + C,
\end{aligned} \tag{18}$$

and

$$\int \frac{dx}{\sinh^{2n} x} = \begin{cases} (-1)^{n+1}\dfrac{(2n-2)!!}{(2n-1)!!}\cosh x \displaystyle\sum_{k=1}^{n} \frac{(2k-3)!!(-1)^k}{(2k-2)!!\sinh^{2k-1} x} + C, & \text{or}\\[2ex] (-1)^{n+1}(n-1)!\displaystyle\sum_{k=1}^{n} \frac{(-1)^k \coth^{2k-1} x}{(2k-1)(k-1)!(n-k)!} + C, \end{cases} \tag{19}$$

where C is a constant. The double factorial $(\cdot)!! : \{-1,0,1,\ldots\} \to \mathbf{N}$ is defined by

$$n!! := \begin{cases} n\cdot(n-2)\cdots 2 & \text{if } n \text{ even} \geq 2,\\ n\cdot(n-2)\cdots 1 & \text{if } n \text{ odd} \geq 1,\\ 1 & \text{if } n \in \{-1,0\}. \end{cases}$$

Note that $(2n)!! = 2^n n!$ for $n \in \mathbf{N}_0$. The finite summation formulae for $\mathfrak{I}_m(x)$ all follow trivially by induction using (17) and the binomial expansion (cf. (1.2.2) in Olver *et al.* (2010) [21])

$$(1 - \coth^2 x)^n = n! \sum_{k=0}^{n} \frac{(-1)^k \coth^{2k} x}{k!(n-k)!}.$$

The formulae (18) and (19) are essentially equivalent to (2.416.2–3) in Gradshteyn & Ryzhik (2007), except (2.416.3) is not defined for the integrand $1/\sinh x$. By applying the limits of integration from the definition of $\mathcal{I}_d(\rho)$ in (15) to (18) and (19) we obtain the following finite summation expressions for $\mathcal{I}_d(\rho)$

$$\mathcal{I}_d(\rho) = \begin{cases} (-1)^{d/2-1} \dfrac{(d-3)!!}{(d-2)!!} \left[\log \coth \dfrac{\rho}{2} + \cosh \rho \displaystyle\sum_{k=1}^{d/2-1} \dfrac{(2k-2)!!(-1)^k}{(2k-1)!! \sinh^{2k} \rho} \right] & \text{if } d \text{ even}, \\ \left\{ \begin{array}{l} (-1)^{(d-1)/2} \left[\dfrac{(d-3)!!}{(d-2)!!} \right. \\ \qquad \left. + \left(\dfrac{d-3}{2} \right)! \displaystyle\sum_{k=1}^{(d-1)/2} \dfrac{(-1)^k \coth^{2k-1} \rho}{(2k-1)(k-1)!((d-2k-1)/2)!} \right], \\ \text{or} \\ (-1)^{(d-1)/2} \dfrac{(d-3)!!}{(d-2)!!} \left[1 + \cosh \rho \displaystyle\sum_{k=1}^{(d-1)/2} \dfrac{(2k-3)!!(-1)^k}{(2k-2)!! \sinh^{2k-1} \rho} \right], \end{array} \right\} & \text{if } d \text{ odd}. \end{cases} \tag{20}$$

Moreover, the antiderivative (indefinite integral) can be given in terms of the Gauss hypergeometric function

$$\int \frac{d\rho}{\sinh^{d-1} \rho} = \frac{-1}{(d-1) \cosh^{d-1} \rho} \, {}_2F_1 \left(\frac{d-1}{2}, \frac{d}{2}; \frac{d+1}{2}; \frac{1}{\cosh^2 \rho} \right) + C, \tag{21}$$

where $C \in \mathbf{R}$. The Gauss hypergeometric function ${}_2F_1 : \mathbf{C}^2 \times (\mathbf{C} \setminus -\mathbf{N}_0) \times \{z \in \mathbf{C} : |z| < 1\} \to \mathbf{C}$ can be defined in terms of the infinite series

$${}_2F_1(a, b; c; z) := \sum_{n=0}^{\infty} \frac{(a)_n (b)_n}{(c)_n n!} z^n$$

(see (15.2.1) in Olver *et al.* (2010) [21]), and elsewhere in z by analytic continuation. (see (2.1.5) in Andrews, Askey & Roy 1999), The Pochhammer symbol (rising factorial) $(\cdot)_l : \mathbf{C} \to \mathbf{C}$ is defined by

$$(z)_n := \prod_{i=1}^{n} (z + i - 1),$$

where $l \in \mathbf{N}_0$. Note that

$$(z)_l = \frac{\Gamma(z+l)}{\Gamma(z)},$$

for all $z \in \mathbf{C} \setminus -\mathbf{N}_0$. The gamma function $\Gamma : \mathbf{C} \setminus -\mathbf{N}_0 \to \mathbf{C}$ (see Chapter 5 in Olver *et al.* (2010) [21]), which is ubiquitous in special function theory, is an important combinatoric function which generalizes the factorial function over the natural numbers. It is naturally defined over the right-half complex plane through Euler's integral (see (5.2.1) in Olver *et al.* (2010) [21])

$$\Gamma(z) := \int_0^\infty t^{z-1} e^{-t} dt,$$

$\operatorname{Re} z > 0$. Some properties of the gamma function, which we will find useful are included below. An important formula which the gamma function satisfies is the duplication formula (i.e., (5.5.5) in Olver *et al.* (2010) [21])

$$\Gamma(2z) = \frac{2^{2z-1}}{\sqrt{\pi}} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right), \tag{22}$$

provided $2z \notin -\mathbf{N}_0$,

The antiderivative (21) is verified as follows. By using

$$\frac{d}{dz} {}_2F_1(a, b; c; z) = \frac{ab}{c} {}_2F_1(a+1, b+1; c+1; z)$$

(see (15.5.1) in Olver *et al.* (2010) [21]), and the chain rule, we can show that

$$\begin{aligned} \frac{d}{d\rho} \frac{-1}{(d-1)\cosh^{d-1}\rho} {}_2F_1\left(\frac{d-1}{2}, \frac{d}{2}; \frac{d+1}{2}; \frac{1}{\cosh^2\rho}\right) = \\ \frac{\sinh\rho}{\cosh^d\rho} {}_2F_1\left(\frac{d-1}{2}, \frac{d}{2}; \frac{d+1}{2}; \frac{1}{\cosh^2\rho}\right) \\ + \frac{d\sinh\rho}{(d+1)\cosh^{d+2}\rho} {}_2F_1\left(\frac{d+1}{2}, \frac{d+2}{2}; \frac{d+3}{2}; \frac{1}{\cosh^2\rho}\right). \end{aligned}$$

The second hypergeometric function can be simplified using Gauss' relations for contiguous hypergeometric functions, namely

$$z\, {}_2F_1(a+1, b+1; c+1; z) = \frac{c}{a-b}[{}_2F_1(a, b+1; c; z) - {}_2F_1(a+1, b; c; z)]$$

(see p. 58 in Erdélyi *et al.* (1981) [6]), and

$${}_2F_1(a, b+1; c; z) = \frac{b-a}{b} {}_2F_1(a, b; c; z) + \frac{a}{b} {}_2F_1(a+1, b; c; z)$$

(see (15.5.12) in Olver *et al.* (2010) [21]). By doing this, the term with the hypergeometric function cancels leaving only a term which is proportional to a binomial through

$${}_2F_1(a, b; b; z) = (1-z)^{-a}$$

(see (15.4.6) in Olver *et al.* (2010) [21]), which reduces to $1/\sinh^{d-1}\rho$. By applying the limits of integration from the definition of $\mathcal{I}_d(\rho)$ in (15) to (21) we obtain the following Gauss hypergeometric representation

$$\mathcal{I}_d(\rho) = \frac{1}{(d-1)\cosh^{d-1}\rho} {}_2F_1\left(\frac{d-1}{2}, \frac{d}{2}; \frac{d+1}{2}; \frac{1}{\cosh^2\rho}\right). \tag{23}$$

Using (23), we can write another expression for $\mathcal{I}_d(\rho)$. Applying Eulers's transformation

$$ {}_2F_1(a,b;c;z) = (1-z)^{c-a-b}\,{}_2F_1\left(c-a,c-b;c;z\right) $$

(see (2.2.7) in Andrews, Askey & Roy (1999) [2]), to (23) produces

$$ \mathcal{I}_d(\rho) = \frac{1}{(d-1)\cosh\rho\,\sinh^{d-2}\rho}\,{}_2F_1\left(\frac{1}{2},1;\frac{d+1}{2};\frac{1}{\cosh^2\rho}\right). $$

Our derivation for a fundamental solution of Laplace's equation on the R-radius hyperboloid $\mathbf{H}^d_R$ in terms of the associated Legendre function of the second kind is as follows. By starting with (23) and the definition of the associated Legendre function of the second kind $Q^\mu_\nu : (1,\infty) \to \mathbf{C}$, namely

$$ Q^\mu_\nu(z) := \frac{\sqrt{\pi}e^{i\pi\mu}\Gamma(\nu+\mu+1)(z^2-1)^{\mu/2}}{2^{\nu+1}\Gamma(\nu+\frac{3}{2})z^{\nu+\mu+1}}\,{}_2F_1\left(\frac{\nu+\mu+2}{2},\frac{\nu+\mu+1}{2};\nu+\frac{3}{2};\frac{1}{z^2}\right), $$

for $|z|>1$ and $\nu+\mu+1 \notin -\mathbf{N}_0$ (see (8.1.3) in Abramowitz & Stegun (1972) [1]), we derive

$$ {}_2F_1\left(\frac{d-1}{2},\frac{d}{2};\frac{d+1}{2};\frac{1}{\cosh^2\rho}\right) = \frac{2^{d/2}\Gamma\left(\frac{d+1}{2}\right)\cosh^{d-1}\rho}{\sqrt{\pi}e^{i\pi(d/2-1)}(d-2)!\sinh^{d/2-1}\rho}Q^{d/2-1}_{d/2-1}(\cosh\rho). \qquad (24) $$

We have therefore verified that the harmonics computed in section 3.1, namely $u^{d,0}_{2+}$, give an alternate form of a fundamental solution for the Laplacian on the hyperboloid. Using the duplication formula for gamma functions (22), (23), and (24), we derive

$$ \mathcal{I}_d(\rho) = \frac{e^{-i\pi(d/2-1)}}{2^{d/2-1}\Gamma\left(d/2\right)\sinh^{d/2-1}\rho}Q^{d/2-1}_{d/2-1}(\cosh\rho). $$

Notice that our chosen fundamental solutions of the Laplacian on the hyperboloid have the property that they tend towards zero at infinity (even for the $d=2$ case, unlike Euclidean fundamental solutions of the Laplacian). Therefore these Green's functions are positive (see Grigor'yan (1983) [10]; Grigor'yan (1985) [11]) and hence $\mathbf{H}^d_R$ is not parabolic. Note that as a result of our proof, we see that the relevant associated Legendre functions of the second kind for $d \in \{2,3,4,5,6,7\}$ are (cf. (20))

$$ \begin{aligned}
&Q_0(\cosh\rho) = \log\coth\frac{\rho}{2},\\
&\frac{1}{\sinh^{1/2}\rho}Q^{1/2}_{1/2}(\cosh\rho) = i\sqrt{\frac{\pi}{2}}(\coth\rho-1),\\
&\frac{1}{\sinh\rho}Q^1_1(\cosh\rho) = \log\coth\frac{\rho}{2} - \frac{\cosh\rho}{\sinh^2\rho},\\
&\frac{1}{\sinh^{3/2}\rho}Q^{3/2}_{3/2}(\cosh\rho) = 3i\sqrt{\frac{\pi}{2}}\left(-\frac{1}{3}\coth^3\rho+\coth\rho-\frac{2}{3}\right),\\
&\frac{1}{\sinh^2\rho}Q^2_2(\cosh\rho) = 3\,\log\coth\frac{\rho}{2} - 2\frac{\cosh\rho}{\sinh^4\rho} - 3\frac{\cosh\rho}{\sinh^2\rho}, \quad \text{and}\\
&\frac{1}{\sinh^{5/2}\rho}Q^{5/2}_{5/2}(\cosh\rho) = 15i\sqrt{\frac{\pi}{2}}\left(\frac{1}{15}\coth^5\rho-\frac{2}{3}\coth^3\rho+\coth\rho-\frac{8}{15}\right).
\end{aligned} $$

The constant c_0 in a fundamental solution for the Laplace operator on the hyperboloid (16) is computed by locally matching up the singularity to a fundamental solution for the Laplace operator in Euclidean space, Theorem 3.2. The coefficient c_0 depends on d. It is determined as follows. For $d \geq 3$ we take the asymptotic expansion for $c_0\mathcal{I}_d(\rho)$ as ρ approaches zero and match this to a fundamental solution of Laplace's equation for Euclidean space given in Theorem 3.2. This yields

$$c_0 = \frac{\Gamma\left(d/2\right)}{2\pi^{d/2}}. \tag{25}$$

For $d = 2$ we take the asymptotic expansion for

$$c_0\mathcal{I}_2(\rho) = c_0 \log \coth \frac{\rho}{2} \simeq c_0 \log \|\mathbf{x} - \mathbf{x}'\|^{-1}$$

as ρ approaches zero, and match this to $\mathcal{G}^2(\mathbf{x}, \mathbf{x}') = (2\pi)^{-1} \log \|\mathbf{x} - \mathbf{x}'\|^{-1}$, therefore $c_0 = (2\pi)^{-1}$. This exactly matches (25) for $d = 2$. The derivation that $\mathcal{I}_d(\rho)$ is an fundamental solution of the Laplace operator on the hyperboloid $\mathbf{H}^d_R$ and the functions for $\mathcal{I}_d(\rho)$ are computed above.

The sectional curvature of a pseudo-sphere of radius R is $-1/R^2$. Hence using results in Losev (1986) [19], all equivalent expressions in Theorem 3.1 can be used for a fundamental solution of the Laplace-Beltrami operator on the R-radius hyperboloid $\mathbf{H}^d_R$ (cf. section 2), namely (where R is now a free parameter)

$$\mathcal{H}^d_R(\mathbf{x}, \mathbf{x}') := \frac{\Gamma\left(d/2\right)}{2\pi^{d/2}R^{d-2}}\mathcal{I}_d\left(\rho\right).$$

The proof of Theorem 3.1 is complete.

Furthermore, due to a theorem proved in [19], all equivalent expressions for $\mathcal{I}_d(\rho)$ in Theorem 3.1 represent upper bounds for a fundamental solution of the Laplace-Beltrami operator on non-compact Riemannian manifolds with negative sectional curvature not exceeding $-1/R^2$ with $R > 0$.

We would also like to mention that a similar computation for a fundamental solution of Laplace's equation on the positive-constant sectional curvature compact manifold, the R-radius hypersphere, has recently been computed in Cohl (2011) [5].

4. Uniqueness of fundamental solution in terms of decay at infinity

It is clear that in general a fundamental solution of Laplace's equation in the hyperboloid model of hyperbolic geometry $\mathcal{H}^d_R$ is not unique since one can add any harmonic function $h : \mathbf{H}^d_R \to \mathbf{R}$ to $\mathcal{H}^d_R$ and still obtain a solution to

$$-\Delta\mathcal{H}^d_R(\mathbf{x}, \mathbf{x}') = \delta_g(\mathbf{x}, \mathbf{x}'),$$

since h is in the kernel of $-\Delta$.

Proposition 4.1. *There exists precisely one C^∞-function $H : (\mathbf{H}^d_R \times \mathbf{H}^d_R) \setminus \{(\mathbf{x}, \mathbf{x}) : \mathbf{x} \in \mathbf{H}^d_R\} \to \mathbf{R}$ such that for all $\mathbf{x}' \in \mathbf{H}^d_R$ the function $H_{\mathbf{x}'} : \mathbf{H}^d_R \setminus \{\mathbf{x}'\} \to \mathbf{R}$ defined by $H_{\mathbf{x}'}(\mathbf{x}) := H(\mathbf{x}, \mathbf{x}')$ is a distribution on $\mathbf{H}^d_R$ with*

$$-\Delta H_{\mathbf{x}'} = \delta_g(\cdot, \mathbf{x}')$$

and

$$\lim_{d(\mathbf{x},\mathbf{x}')\to\infty} H_{\mathbf{x}'}(\mathbf{x}) = 0, \tag{26}$$

where $d(\mathbf{x}, \mathbf{x}')$ is the geodesic distance between two points $\mathbf{x}, \mathbf{x}' \in \mathbf{H}^d_R$.

Proof. Existence: clear. Uniqueness. Suppose H and $\tilde{H}$ are two such functions. Let $\mathbf{x}' \in \mathbf{H}^d_R$. Define the C^∞-function $h : \mathbf{H}^d_R \setminus \{\mathbf{x}'\} \to \mathbf{R}$ by $h = H_{\mathbf{x}'} - \tilde{H}_{\mathbf{x}'}$. Then h is a distribution on $\mathbf{H}^d_R$ with $-\Delta h = 0$. Since $\mathbf{H}^d_R$ is locally Euclidean one has by local elliptic regularity that h can be extended to a C^∞-function $\hat{h} : \mathbf{H}^d_R \to \mathbf{R}$. It follows from (26) for H and $\tilde{H}$ that

$$\lim_{d(\mathbf{x},\mathbf{x}')\to\infty} \hat{h}(\mathbf{x}) = 0. \tag{27}$$

The strong elliptic maximum/minimum principle on a Riemannian manifold for a bounded domain Ω states that if u is harmonic, then the supremum/infimum of u in Ω coincides with the supremum/infimum of u on the boundary $\partial\Omega$. By using a compact exhaustion sequence Ω_k in a non-compact connected Riemannian manifold and passing to a subsequence $\mathbf{x}_k \in \partial\Omega_k$ such that $\mathbf{x}_k \to \infty$, the strong elliptic maximum/minimum principle can be extended to non-compact connected Riemannian manifolds with boundary conditions at infinity (see for instance section 8.3.2 in Grigor'yan (2009) [12]). Taking $\Omega_k \subset \mathbf{H}^d_R$, the strong elliptic maximum/minimum principle for non-compact connected Riemannian manifolds implies using (27) that $\hat{h} = 0$. Therefore $h = 0$ and $H(\mathbf{x}, \mathbf{x}') = \tilde{H}(\mathbf{x}, \mathbf{x}')$ for all $\mathbf{x} \in \mathbf{H}^d_R \setminus \{\mathbf{x}'\}$.

By Proposition 4.1, for $d \geq 2$, the function $\mathcal{H}^d_R$ is the unique normalized fundamental solution of Laplace's equation which satisfies the vanishing decay (26).

Acknowledgements

Much thanks to Simon Marshall, A. Rod Gover, Tom ter Elst, Shaun Cooper, George Pogosyan, Willard Miller, Jr., and Alexander Grigor'yan for valuable discussions. I would like to express my gratitude to Carlos Criado Cambón in the Facultad de Ciencias at Universidad de Málaga for his assistance in describing the global geodesic distance function in the hyperboloid model. We would also like to acknowledge two anonymous referees whose comments helped improve this paper. I acknowledge funding for time to write this paper from the Dean of the Faculty of Science at the University of Auckland in the form of a three month stipend to enhance University of Auckland 2012 PBRF Performance. Part of this work was conducted while H. S. Cohl was a National Research

Council Research Postdoctoral Associate in the Information Technology Laboratory at the National Institute of Standards and Technology, Gaithersburg, Maryland, U.S.A.

References

[1] M. Abramowitz and I. A. Stegun. *Handbook of mathematical functions with formulas, graphs, and mathematical tables*, volume 55 of *National Bureau of Standards Applied Mathematics Series*. U.S. Government Printing Office, Washington, D.C., 1972.

[2] G. E. Andrews, R. Askey, and R. Roy. *Special functions*, volume 71 of *Encyclopedia of Mathematics and its Applications*. Cambridge University Press, Cambridge, 1999.

[3] E. Beltrami. Essai d'interprétation de la géométrie noneuclidéenne. Trad. par J. Hoüel. *Annales Scientifiques de l'École Normale Supérieure*, 6:251–288, 1869.

[4] L. Bers, F. John, and M. Schechter. *Partial differential equations*. Interscience Publishers, New York, N.Y., 1964.

[5] H. S. Cohl. Fundamental solution of laplace's equation in hyperspherical geometry. *Symmetry, Integrability and Geometry: Methods and Applications*, 7(108), 2011.

[6] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Higher transcendental functions. Vol. I.* Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981.

[7] A. Erdélyi, W. Magnus, F. Oberhettinger, and F. G. Tricomi. *Higher transcendental functions. Vol. II.* Robert E. Krieger Publishing Co. Inc., Melbourne, Fla., 1981.

[8] G. B. Folland. *Introduction to partial differential equations*. Number 17 in Mathematical Notes. Princeton University Press, Princeton, 1976.

[9] D. Gilbarg and N. S. Trudinger. *Elliptic partial differential equations of second order*. Number 224 in Grundlehren der mathematischen Wissenschaften. Springer-Verlag, Berlin etc., second edition, 1983.

[10] A. A. Grigor'yan. Existence of the Green function on a manifold. *Russian Mathematical Surveys*, 38(1(229)):161–162, 1983.

[11] A. A. Grigor'yan. The existence of positive fundamental solutions of the Laplace equation on Riemannian manifolds. *Matematicheskie Zametki*, 128(170)(3):354–363, 446, 1985.

[12] A. A. Grigor'yan. *Heat kernel and analysis on manifolds*, volume 47 of *AMS/IP Studies in Advanced Mathematics*. American Mathematical Society, Providence, RI, 2009.

[13] C. Grosche, G. S. Pogosyan, and A. N. Sissakian. Path-integral approach for superintegrable potentials on the three-dimensional hyperboloid. *Physics of Particles and Nuclei*, 28(5):486–519, 1997.

[14] S. Helgason. *Groups and geometric analysis: Integral geometry, invariant differential operators, and spherical functions*, volume 113 of *Pure and Applied Mathematics*. Academic Press Inc., Orlando, FL, 1984.

[15] L. Hostler. Vector spherical harmonics of the unit hyperboloid in Minkowski space. *Journal of Mathematical Physics*, 18(12):2296–2307, 1977.

[16] A. A. Izmest'ev, G. S. Pogosyan, A. N. Sissakian, and P. Winternitz. Contractions of Lie algebras and separation of variables. The n-dimensional sphere. *Journal of Mathematical Physics*, 40(3):1549–1573, 1999.

[17] A. A. Izmest'ev, G. S. Pogosyan, A. N. Sissakian, and P. Winternitz. Contractions of Lie algebras and the separation of variables: interbase expansions. *Journal of Physics A: Mathematical and General*, 34(3):521–554, 2001.

[18] J. M. Lee. *Riemannian manifolds*, volume 176 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1997.

[19] A. G. Losev. Harmonic functions on manifolds of negative curvature. *Matematicheskie Zametki*, 40(6):738–742, 829, 1986.

[20] M. N. Olevskiĭ. Triorthogonal systems in spaces of constant curvature in which the equation

$\Delta_2 u + \lambda u = 0$ allows a complete separation of variables. *Matematicheskiĭ Sbornik*, 27(69):379–426, 1950. (in Russian).

[21] F. W. J. Olver, D. W. Lozier, R. F. Boisvert, and C. W. Clark, editors. *NIST handbook of mathematical functions*. Cambridge University Press, Cambridge, 2010.

[22] G. S. Pogosyan and P. Winternitz. Separation of variables and subgroup bases on n-dimensional hyperboloids. *Journal of Mathematical Physics*, 43(6):3387–3410, 2002.

[23] W. P. Thurston. *Three-dimensional geometry and topology. Vol. 1*, volume 35 of *Princeton Mathematical Series*. Princeton University Press, Princeton, NJ, 1997. Edited by Silvio Levy.

[24] R. J. Trudeau. *The non-Euclidean revolution*. Birkhäuser, Boston, 1987.

[25] N. Ja. Vilenkin. *Special functions and the theory of group representations*. Translated from the Russian by V. N. Singh. Translations of Mathematical Monographs, Vol. 22. American Mathematical Society, Providence, R. I., 1968.